

Improving physical abilities and the most efficient methods in adults over 40 years old.

Kristian Andrea³, Anesti Qeleshi², Redon Zotaj³

¹Sports University of Tirana, Faculty of Physical Activity and Recreation

²Sports University of Tirana, Faculty of Movement Sciences

Correspondence:(aandrea@ust.edu.al)

Abstract

Only a small percentage of adults engage in regular physical activity, which means a decline in quality of life and a very poor result compared to Europe (Osiński, 2013). The purpose of this literature review is to find the most contemporary methods from the most ranked scientific works on the improvement of physical abilities and the most efficient methods in adults over 40 years old. Digital databases such as Scopus, Medline, JStore, Embase, Cochrane Central, PsycInfo, Scholar, and SPORTDiscus were used in this literature review. Inclusion criteria are: individuals must be healthy and average age at least 40 years old. Many other studies such as (Skelton et al., 1994; Foldvari et al., 2000; Bean et al., 2007) report that physical functions are more related to muscle power than to muscle strength or mass. Strength training has many benefits on physical function in older adults compared to traditional strength training (Tschopp et al., 2011; Da Rosa., 2019). Harmonization or proper use of variables such as exercise selection, breaks, number of sets, number of repetitions, frequency affect the increase in the level of muscular fitness. Strength, muscular endurance, muscular strength and muscular hypertrophy are the characteristics that should be trained (Ratamess et al., 2009). While in an all-inclusive program resistance training improves the cardiovascular system (Fleck. 1988), limits the risk of coronary heart disease (Goldberg. 1989; Hurley & Kokkinos, 1987), helps prevent osteoporosis (Layne & Nelson. 1999), removes the risk of diabetes (Miller et al., 1984), promotes maintaining a healthy weight and losing excess weight (Evans. 1999), reduces the risk of cancer in the spine (Koffler et al., 1992), helps in psychological/mental well-being (Ewart. 1989) as well as improves dynamic stability and maintenance of functional capacity

(Evans, 1999). In conclusion from the study of Balachandran et al. (2022) found that strength training leads to improvements in physical function more than traditional strength training.

Keywords: fitness, aging, power, strength, exercises

Introduction

Only a small percentage of adults engage in regular physical activity, which means a decline in quality of life and a very poor result compared to Europe (Osiński, 2013). The ability to perform activities of daily living (ADL) is vital to living independently. Age-related loss in muscle strength can compromise (Hairi et al., 2010) this ability and lead to disability. For example, the progression of muscle weakness limits the ability to grasp an object which further hinders the ability to open a jar. Experiencing difficulty in ADL and relying on others is not only associated with reduced quality of life (Hellstrom et al., 2004) but also increased likelihood (Gaugler et al., 2007) of long-term nursing (staying at home).

As a result, the population of the third age will dominate, which will be a serious problem for the economy of the state due to poor health. Therefore, maximum efforts are needed to improve the quality of health and life of the third-age population. Szukalski (2011) pointed out that the decrease in income as a result of the aging of the population means the need to increase the professional activity of the elderly. But this need has other disadvantages such as low physical performance as a result of involitional structural changes in the internal organs, the respiratory system, the cardiovascular circulation and the passive and active motoric system (English & Paddon-Jones, 2010).

As a result, there is a need for health promoters, the education system and the media to encourage society about regular health education. When encouraging people to work long hours, it should be accompanied by the right physical condition. This requires a good balance of the scope of professional activity and the absence of people at work. The citizens of the Scandinavian countries and the Netherlands are in the best physical condition (KMPG, 2012) stating that 50-70% of them admit that they perform physical activity at least once a week. This percentage also includes people after the age of 50, noticing lower expenses for the treatment of the elderly. Studies show that the citizens of the Scandinavian countries and the Netherlands are in the best physical condition (KMPG, 2012) stating that 50-70% of them admit that they perform physical activity at least once a week. This percentage also includes people after the age of 50, noticing lower expenses for the treatment of the elderly.

The focus on each training component depends on the tasks each training period has. The realization of the performance has a basis which is physical preparation. The higher the

physical preparation, the more effective the technical, tactical and psychological components will be. The insufficient level of PF (physical preparation) reflects in the use of human potential during sports performance, the technique is broken and fatigue appears quickly. In the training components, physical preparation is like a foundation stone on the basis of which the other components "rise". The time required to create a state of higher adaptation depends on the complexity of the skill and on the athlete's physiological and psychological difficulties. The more difficult and complex the sport, the longer the necessary training period will be to promote neuromuscular adaptation.

Regardless of age, a suitable physical activity will bring about the necessary changes in the body (Wernbom et al., 2007). A suitable physical activity is strength and endurance training. According to forecasts, the number of senior citizens is expected to double by 2050 to approximately 1.5 billion (Werner, 2010). However, there are some intervention programs that help slow down the decline in physical functions. One of these programs is strength training to prevent or delay movement limitations in elderly individuals. According to Chmelo et al., (2015) strength training has been recommended to improve physical functions in elderly individuals. However, it is muscle strength that has emerged as a key factor in physical function. Researcher Reid et al., (2014) prove that it is muscle power that decreases at a faster rate compared to muscle strength during the aging period.

The purpose of this literature review is to find the most contemporary methods from the most ranked scientific works on the improvement of physical abilities and the most efficient methods in adults over 40 years old.

Methods

Digital databases such as Scopus, Medline, JStore, Embase, Cochrane Central, PsycInfo, Scholar, and SPORTDiscus were used in this literature review. There was a restriction on the year of publications (last 10-15 years). Inclusion criteria are: individuals must be healthy and average age at least 40 years old. The resistance training intervention program included machine-related exercises, body weight exercises, ergometric bicycles, elastic belts, etc.

Results

Researcher Reid et al. (2014) prove that it is muscle power that decreases at a faster rate compared to muscle strength during the aging period. Many other studies such as (Skelton et al., 1994; Foldvari et al., 2000; Bean et al., 2007) report that physical functions are more related to muscle power than to muscle strength or mass. Strength training has many benefits on physical function in older adults compared to traditional strength training (Tschopp et al., 2011;

Da Rosa., 2019). These studies include plyometric training with exercises such as (CMJ and depth jump) (Miszko et al., 2003; Ramirez-Campillo et al., 2014). According to Vetrovsky et al. (2019) unlike power training which uses slow controlled eccentricity, plyometric training uses fast eccentricity. Previous studies examining the effects of strength training have used different interventions. As an example, standing strength training exercises were used for the experimental group while seated strength training exercises were used for the control group (Bean et al., 2009). The inclusion of many intervention programs makes it difficult to understand the impact that strength training has on physical function.

Some types of strength training require a special methodology (Stefaniak, 2006; Zajac et al., 2010). Before starting a training program you must apply a preparatory cycle to adapt to the subsequent program. This preparatory cycle can last from several weeks to several months and has a health and self-educational value to encourage people to engage in physical activity. Resistance training induces biochemical changes in skeletal muscle and thus improves whole body functions (Wernbom et al., 2007). Resistance training causes an increase in the activity of collagen proteins, which affects the increase in muscle turnover (Burd et al., 2010). The study by Davis et al. (2008) prove that the combination of strength training and aerobic training for 11 weeks improves endurance and strength. The high-intensity training group changed body composition more compared to the low-intensity training group. Many literatures suggest the use of resistance exercises. In conclusion, the study of Davis et al. (2008) proved that the combination of strength training and aerobic training for 11 weeks improves endurance and strength.

In the study of Marques et al. (2011) resistance training produced positive changes in bone density (in women), fat reduction and improved balance. Also the researcher Borer. (2005) confirm changes in the inhibition of osteoporosis after the application of resistance training. Researchers Willis et al. (2012) and Ormsbee et al. (2009) proved that isometric exercise improves resting metabolism.

The aim of the study of Kalwa et al. (2019) is to find a suitable method to inhibit the involutionary processes as well as to improve the physical abilities of people over 40 years old. Also, this work aims to determine the loads and the dose of physical efforts, which must be looked at carefully to avoid injuries and to eliminate the negative effects of resistance training. 178 adults, 92 of whom were women and 86 men aged 40-61, participated in this study. They participated in an 8-month training program. The participants in the study were divided into two groups - group 1 (51-61 years old) and group 2 (40-50 years old). Before applying the strength program, a preparatory training cycle was carried out. The training was carried out 3

times a week. Two exercises were performed for each major body part. 15 minutes of warm-up and 60-80 minutes of strength training as well as 10 minutes of stretching. The final results from this study show that the exercise program provides beneficial effects in the fight against the symptoms of aging. The data showed changes in body composition and physical fitness in both groups but the group where the higher intensity exercise was applied had greater changes. Improved fitness was seen in both women and men. The greatest changes were observed in physical fitness.

The meta-analysis of the study of Balachandran et al. (2022) evaluated the impact of strength training on physical functions in the third age. A small number of findings showed that strength training improved physical function more than traditional strength training. The above study only evaluated meta-analyses or studies related to strength training without using plyometric exercises. Two other studies were not included in this meta-analysis because the experimental group performed standing strength training exercises while the control group used seated strength training exercises (Bean, 2004).

Also, two other studies were not included in the analysis because they performed CMJ (Miszko et al., 2003), and the second study combined circuit training with strength training (Balachandran et al., 2014).

The study by Balachandran et al. (2022) focused only on strength training without the intervention of other types of training. To assess physical functions, most studies used the GUG test and the chair sitting test. These two above mentioned tests are used in a fast rhythm while the I Timed up and go test is used in a regular rhythm. For the GUG test we observed a mean improvement of 1.85 seconds SD 0.62 while for the chair stand test a greater improvement was observed with 4.3 seconds SD 0.56. Strength training produces an increase in strength in type II muscle fibers independent of age and gender (Claflin et al., 1985, 2011). According to Lexell (1995) the fibers of the second type have a loss during the aging period. As seen the improvement of strength and power is responsible for the limitation of movements (Santanasto et al., 2017). Consequently, the improvement of muscular strength provides a biological basis for the improvement of physical functions. While walking speed, strength and muscle mass are not related. In many studies, the same levels of intensity and volume were used for both groups, so we do not expect to have different results of strength and muscle mass. While studies that showed improvements in walking speed applied walking as an exercise program (Pahor et al., 2014). Adverse effects were minimal in both groups.

Discussion

It should be noted that some forms of physical activity that may be of particular importance to an aging population (eg, balance training) may not fit this standard definition. This activity can be carried out during leisure or working hours, and surveys of older adults (Harada, 2001) should include both paid and unpaid (volunteer) work. Moderate physical activity does not appear to be sufficient in terms of increasing participation in these activities among older adults (Dunn, Anderson, & Jakicic, 1998). In order to effect long-term changes in behavior, it is necessary to identify, examine and begin to provide information on the barriers to physical activity that affect adults so that they do not become obstacles from the transition from a passive to a physically active one (Dishman & Sallis, 1994). Harmonization or proper use of variables such as exercise selection, breaks, number of sets, number of repetitions, frequency affect the increase in the level of muscular fitness. Strength, muscular endurance, muscular strength and muscular hypertrophy are the characteristics that should be trained (Ratamess et al., 2009). While in an all-inclusive program resistance training improves the cardiovascular system (Fleck. 1988), limits the risk of coronary heart disease (Goldberg. 1989; Hurley & Kokkinos, 1987), helps prevent osteoporosis (Layne & Nelson. 1999), removes the risk of diabetes (Miller et al., 1984), promotes maintaining a healthy weight and losing excess weight (Evans. 1999), reduces the risk of cancer in the spine (Koffler et al., 1992), helps in psychological/mental well-being (Ewart. 1989) as well as improves dynamic stability and maintenance of functional capacity (Evans. 1999). In conclusion from the study of Balachandran et al. (2022) found that strength training leads to improvements in physical function more than traditional strength training. The study evaluated only meta-analyses or studies related to strength training without using plyometric exercises. For the GUG test we observed a mean improvement of 1.85 seconds SD 0.62 while for the chair stand test a greater improvement was observed with 4.3 seconds SD 0.56. Strength training produces an increase in strength in type II muscle fibers independent of age and gender (Claflin et al., 1985). According to Lexell (1995) the fibers of the second type have a loss during the aging period. As seen the improvement of strength and power is responsible for the limitation of movements (Santanasto et al., 2017). Consequently, the improvement of muscular strength provides a biological basis for the improvement of physical functions. According to studies (Fried & Guralnik, 1997; Janssen et al., 2002; Bergen et al., 2014) aging is associated with a decrease in physical functions, loss of physical independence as a consequence and a decrease in the quality of life. The study by Katz et al. (1983) emphasize that the extension of life expectancy must necessarily be accompanied by the good preservation of physical functions. According to

forecasts, the number of senior citizens is expected to double by 2050 to approximately 1.5 billion (Werner, 2010). However, there are some intervention programs that help slow down the decline in physical functions. One of these programs is strength training to prevent or delay movement limitations in elderly individuals. According to Chmelo et al. (2015) strength training has been recommended to improve physical functions in elderly individuals. However, it is muscle strength that has emerged as a key factor in physical function.

References

- Balachandran, AT, Steele, J., Angielczyk, D., Belio, M., Schoenfeld, BJ, Quiles, N., ... & Abou-Setta, AM (2022). Comparison of power training vs traditional strength training on physical function in older adults: a systematic review and meta-analysis. *JAMA network open*, 5 (5), e2211623-e2211623.
- Balachandran, A., Krawczyk, SN, Potiaumpai, M., & Signorile, JF (2014). High-speed circuit training vs hypertrophy training to improve physical function in sarcopenic obese adults: A randomized controlled trial. *Experimental gerontology*, 60, 64-71. doi:10.1016/j.exger.2014.09.016
- Bean, JF, Kiely, DK, LaRose, S., Alian, J., & Frontera, WR (2007). Is stair climb power a clinically relevant measure of leg power impairments in at-risk older adults?. *Archives of physical medicine and rehabilitation*, 88 (5), 604-609. doi:10.1016/j.apmr.2007.02.004
- Bean, JF, Kiely, DK, LaRose, S., O'Neill, E., Goldstein, R., & Frontera, WR (2009). Increased velocity exercise specific to task training versus the National Institute on Aging's strength training program: changes in limb power and mobility. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*, 64 (9), 983-991. doi:10.1093/gerona/glp056
- Bean, JF, Herman, S., Kiely, DK, Frey, IC, Leveille, SG, Fielding, RA, & Frontera, WR (2004). Increased Velocity Exercise Specific to Task (InVEST) training: a pilot study exploring effects on leg power, balance, and mobility in community-dwelling older women. *Journal of the American Geriatrics Society*, 52 (5), 799-804. doi:10.1111/j.1532-5415.2004.52222.x
- Bergen G, Stevens MR, Burns ER. Falls and Fall Injuries Among Adults Aged ≥ 65 Years — United States, 2014. *MMWR Morb Mortal Wkly Rep* 2016;65:993 –998. DOI: [http://dx.doi.org/10.15585/mmwr.mm6537a2external icon](http://dx.doi.org/10.15585/mmwr.mm6537a2external%20icon)

- Borer, KT (2005). Physical activity in the prevention and amelioration of osteoporosis in women: interaction of mechanical, hormonal and dietary factors. *Sports medicine*, 35, 779-830. <https://doi.org/10.2165/00007256-200535090-00004>
- Burd, NA, West, DW, Churchward-Venne, TA, & Mitchell, CJ (2010). Growing collagen, not muscle, with weightlifting and 'growth' hormones. *The Journal of physiology*, 588 (Pt 3), 395. <http://doi.org/10.1113/jphysiol.2009.185306>
- Clafin, DR, Larkin, LM, Cederna, PS, Horowitz, JF, Alexander, NB, Cole, NM, ... & Ashton-Miller, JA (2011). Effects of high- and low-velocity resistance training on the contractile properties of skeletal muscle fibers from young and older humans. *Journal of applied physiology*, 111 (4), 1021-1030
- Chmelo, EA, Crotts, CI, Newman, JC, Brinkley, TE, Lyles, MF, Leng, X., ... & Nicklas, BJ (2015). Heterogeneity of physical function responses to exercise training in older adults. *Journal of the American Geriatrics Society*, 63 (3), 462-469. doi:10.1111/jgs.13322
- Davis, WJ, Wood, DT, Andrews, RG, Elkind, LM, & Davis, WB (2008). Concurrent training enhances athletes' strength, muscle endurance, and other measures. *The Journal of Strength & Conditioning Research*, 22 (5), 1487-1502.
<https://doi.org/10.1519/JSC.0b013e3181739f08>
- da Rosa Orssatto, LB, de la Rocha Freitas, C., Shield, AJ, Pinto, RS, & Trajano, GS (2019). Effects of resistance training concentric velocity on older adults' functional capacity: A systematic review and meta-analysis of randomized trials. *Experimental Gerontology*, 127, 110731.
doi:10.1016/j.exger.2019.110731
- Dishman, RK, & Sallis, JF (1994). Determinants and interventions for physical activity and exercise. In C. Bouchard, RJ Shephard, & T. Stevens (Eds.), *Physical activity, fitness, and health* (pp. 214-238). Champaign, IL: Human Kinetics.
- Dunn, AL, Anderson, RE, & Jakicic, JM (1998). Lifestyle physical activity interventions. *American Journal of Preventive Medicine*, 15(4), 398-412.
- English, KL, & Paddon-Jones, D. (2010). Protecting muscle mass and function in older adults during bed rest. *Current opinion in clinical nutrition and metabolic care*, 13 (1), 34. <http://doi.org/10.1097/MCO.0b013e328333aa66>
- Evans WJ (1999). Exercise training guidelines for the elderly. *Medicine and science in sports and exercise*, 31 (1), 12-17. <https://doi.org/10.1097/00005768-199901000-00004>
- Fleck, SJ (1988). Cardiovascular adaptations to resistance training. *Medicine and science in sports and exercise*, 20 5 Suppl, S146-51 .

- Foldvari, M., Clark, M., Laviolette, LC, Bernstein, MA, Kaliton, D., Castaneda, C., ... & Singh, MAF (2000). Association of muscle power with functional status in community-dwelling elderly women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 55 (4), M192-M199.
doi:10.1093/gerona/55.4.M192
- Fried, LP, & Guralnik, JM (1997). Disability in older adults: evidence regarding significance, etiology, and risk. *Journal of the American Geriatrics Society*, 45 (1), 92-100.
doi:10.1111/j.1532- 5415.1997.tb 00986.x
- Harada, ND, Chiu, V., King, AC, & Stewart, AL (2001). An evaluation of three self-report physical activity instruments for older adults. *Medicine & Science in Sports & Exercise*, 33 (6), 962-970.
- Hurley, BF, & Kokkinos, PF (1987). Effects of weight training on risk factors for coronary artery disease. *Sports Medicine*, 4, 231-238.
- Hairi NN, Cumming RG, Naganathan V, Handelsman DJ, Le Couteur DG, Creasey H et al (2010) Loss of muscle strength, mass (sarcopenia), and quality (specific strength) and its relationship with functional limitation and physical disability: The Concord Health and Aging in Men project. *J Am Geriatr Soc* 58:2055–2062
- Hellström Y, Persson G, Hallberg IR (2004) Quality of life and symptoms among older people living at home. *J Adv Nurs* 48:584– 593
- Gaugler J, Duval S, Anderson K, Kane R (2007) Predicting nursing home admissions in the US: a meta-analysis. *BMC Geriatrics* 7:13
- Goldberg, AP (1989). Aerobic and resistive exercise modify risk factors for coronary heart disease. *Medicine and Science in Sports and Exercise*, 21 (6), 669-674.
- Janssen, I., Heymsfield, SB, & Ross, R. (2002). Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *Journal of the american geriatrics society*, 50 (5), 889-896. doi:10.1046/j.1532-5415.2002.50216.x
- Kalwa, M., Stefaniak, T., Harmacinski, D., & Groffik, B. (2019). A preparation cycle in strength training as a form of health training for adults. *Baltic Journal of Health and Physical Activity*, 11 (1), 7.
- Katz S, Branch LG, Branson MH, Papsidero JA, Beck JC, Greer DS. (1983). Active life expectancy. *N Engl J Med* . 309(20):1218-1224. doi:10.1056/NEJM198311173092005

- Koffler, KH, Menkes, A., Redmond, RA, Whitehead, WE, Pratley, RE, & Hurley, BF (1992). Strength training accelerates gastrointestinal transit in middle-aged and older men. *Medicine and science in sports and exercise*, 24 (4), 415-419
- KMPG. (2012). Analysis of socio-demographic changes and the impact of poor nutrition, insufficient physical activity, addictions, and other risk factors on the prevalence and costs of diabetes and cardiovascular disease in Poland. Current status and forecast 2030. KEMP
- Lexell J, Lexell J. (1995). Human aging, muscle mass, and fiber type composition. *J Gerontol A Biol Sci Med Sci* .50(Spec No):11-16.
- Layne, JE, & Nelson, ME (1999). The effects of progressive resistance training on bone density: a review. *Medicine and science in sports and exercise*, 31 (1), 25-30.
- Miszko, TA, Cress, ME, Slade, JM, Covey, CJ, Agrawal, SK, & Doerr, CE (2003). Effect of strength and power training on physical function in community-dwelling older adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 58 (2), M171-M175. doi:10.1093/gerona/58.2.M171
- Miller, WJ, Sherman, WM, & Ivy, JL (1984). Effect of strength training on glucose tolerance and post-glucose insulin response. *Medicine and science in sports and exercise*, 16 (6), 539-543
- Marques, EA, Mota, J., Machado, L., Sousa, F., Coelho, M., Moreira, P., & Carvalho, J. (2011). Multicomponent training program with weight-bearing exercises elicits favorable bone density, muscle strength, and balance adaptations in older women. *Calcified tissue international*, 88, 117-129. <https://doi.org/10.1007/s00223-010-9437-1>
- Osiński W. (2013). The idea of freedom - against physical culture. *Wych Fiz i Zdrowotne* . 2:4-7.
- Ormsbee, MJ, Choi, MD, Medlin, JK, Geyer, GH, Trantham, LH, Dubis, GS, & Hickner, RC (2009). Regulation of fat metabolism during resistance exercise in sedentary lean and obese men. *Journal of Applied Physiology*, 106 (5), 1529-1537. (Ratamess et al., 2009).
- Pahor, M., Guralnik, JM, Ambrosius, WT, Blair, S., Bonds, DE, Church, TS, ... & LIFE Study Investigators. (2014). Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *Jama*, 311 (23), 2387-2396. doi:10.1001/jama.2014.5616
- Reid, KF, Pasha, E., Doros, G., Clark, DJ, Patten, C., Phillips, EM, ... & Fielding, RA (2014). Longitudinal decline of lower extremity muscle power in healthy and mobility-limited older adults: influence of muscle mass, strength, composition, neuromuscular activation

- and single fiber contractile properties. *European journal of applied physiology*, 114, 29-39. doi:10.1007/s00421-013-2728-2
- Ramírez-Campillo, R., Castillo, A., de la Fuente, CI, Campos-Jara, C., Andrade, DC, Álvarez, C., ... & Izquierdo, M. (2014). High-speed resistance training is more effective than low-speed resistance training to increase functional capacity and muscle performance in older women. *Experimental gerontology*, 58, 51-57. doi:10.1016/j.exger .2014.07.001
- Skelton, DA, Greig, CA, Davies, JM, & Young, A. (1994). Strength, power and related functional ability of healthy people aged 65–89 years. *Age and aging*, 23 (5), 371-377. doi:10.1093/ageing/23.5.371
- Santanasto, AJ, Glynn, NW, Lovato, LC, Blair, SN, Fielding, RA, Gill, TM, ... & LIFE Study Group. (2017). Effect of physical activity versus health education on physical function, grip strength and mobility. *Journal of the American Geriatrics Society*, 65 (7), 1427-1433. doi:10.1111/jgs.14804
- Stefaniak, T., Witkowski, K., & Burdzielowska, M. (2006). Evaluation of sensorimotor efficiency in persons at involution age systematically practicing strength exercises. *Medycyna Sportowa*, 22 (6), 333-340.
- Szukalski S. (2011). Prospects for the development of services in Poland in the light of demographic contraction and aging population. *Soc Unequal Econ Growth* . 18:409-419.
- Tschopp, M., Sattelmayer, MK, & Hilfiker, R. (2011). Is power training or conventional resistance training better for function in elderly persons? A meta-analysis. *Age and aging*, 40 (5), 549-556. doi:10.1093/ageing/afr005
- Vetrovsky, T., Steffl, M., Stastny, P., & Tufano, JJ (2019). The efficacy and safety of lower-limb plyometric training in older adults: a systematic review. *Sports Medicine*, 49, 113-131. doi:10.1007/s40279-018-1018-x
- Zajac A, Wilk M, Poprzęcki S, et al. (2010). Współczesny trening szyły miyówej [Contemporary muscular strength training]. Katowice: AWF; Polish.
- Werner C. The older population: 2010. November 2011. Accessed January 1, 2015. <https://www.census.gov/content/dam/Census/library/publications/2011/dec/c2010br-09.pdf>
- Wernbom, M., Augustsson, J., & Thomeé, R. (2007). The influence of frequency, intensity, volume and mode of strength training on whole muscle cross-sectional area in humans. *Sports medicine*, 37, 225-264. <http://doi.org/10.2165/00007256-200737030-00004>

Willis, LH, Slentz, CA, Bateman, LA, Shields, AT, Piner, LW, Bales, CW, ... & Kraus, WE (2012). Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *Journal of applied physiology*. 2012;113:1831 -1837. <https://doi.org/10.1152/jappphysiol.01370.2011>