

## ARTICLE

### FATTY ACIDS PROFILE OF SOME COMMERCIALY FISH ROE

Carmen POPA <sup>1</sup>, Iulia TIMOFTE <sup>2</sup>, Lucian COPOLOVICI <sup>2</sup>, Dana Maria COPOLOVICI <sup>2\*</sup>

<sup>1</sup> University of Oradea, Biomedical Sciences Doctoral School, 1 University St., 410087, Oradea, Romania

<sup>2</sup> Aurel Vlaicu University, Faculty of Food Engineering, Tourism and Environmental Protection; Institute for Research, Development and Innovation in Technical and Natural Sciences, Romania, 2 Elena Dragoi St., Arad, 310330, Romania

\*Corresponding author: dana.copolovici@uav.ro

**Abstract:** The demand for natural food sources and food ingredients is increasing, and fish roes are known to present high levels of  $\omega$ -3 fatty acids (FA). In this work, we used gas-chromatography and UV spectroscopy (GC-DAD) to characterize the FA profile of six different commercial fish roes. The increase of concentration of  $\omega$ -3 FA is tarama < hering < pike < cod < salmon < lumpfish. Altogether, the fish roes are an important source of unsaturated fatty acids (minimum 26.62% for pike roe) and present a modest source of saturated fatty acids (less than 46.45% for pike roe). Overall, this study provides a preliminary screening of FA profiles between several commercial fish roes.

**Keywords:** fish roe, fatty acids, gas chromatography, mass spectrometry.

## Introduction

The beneficial effects of fish and fish oil consumption on human health are widely documented in the scientific literature (Abdelhamid et al., 2020; Abdelhamid et al., 2018; Alagawany et al., 2019; Cretton et al.). There have been many studies that have verified a reduction in the incidence of a variety of diseases, including cardiovascular disease (de la Guia-galipienso et al., 2021), cancer (de Castro et al., 2022; Franchi et al., 2022; Hsieh et al., 2022; Jiang et al., 2021; Laumann et al., 2022; Stasiewicz et al., 2022; Wang et al., 2022; Zhang et al., 2022), inflammatory (Dawczynski et al., 2018) and autoimmune disorders (AlAmmar et al., 2021; Lofvenborg et al., 2014), as well as psychiatric and mental ailments (Berger et al., 2017; Hansen et al., 2020; Sikka et al., 2021; Yonezawa et al., 2020). Fish liver oil, mushrooms, and other sources of vitamins are the best sources of vitamin D, besides exposure to the sun, for the average person. 1,25-dihydroxyvitamin D is primarily responsible for maintaining calcium and phosphorus balance in the body (de la Guia-galipienso et al., 2021). Many extra-skeletal effects of vitamin D, including those on the immune and cardiovascular (CV) systems, can be detected in human cells and tissues that contain vitamin

D receptors. Blood pressure is regulated by vitamin D's effect on endothelium and smooth muscle cells. This vitamin deficiency has been linked to an increased risk of cardiovascular disease (CVD) death and incidence (de la Guia-galipienso et al., 2021).

Health advantages of a diet containing fish products are attributed to polyunsaturated fatty acids (PUFA), mainly omega-3 fatty acids; e.g., eicosapentaenoic acid: EPA and docosahexaenoic acid: DHA.

The fatty acids (FA) present in four lipid fractions [polar lipids (PL), diacylglycerols (DAG), free fatty acids (FFA), and triacylglycerols (TAG)] of common carp were investigated by Chvalova, D. and J. Spicka in (Chvalova and Spicka, 2016). There were differences across the examined tissues in terms of the amounts of lipid fractions and fatty acids that were classified as: saturated fatty acids (SFA), monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA), and furan fatty acids (F-acids). Forty-nine compounds were identified in this research, from which eight F-acids were discovered in the tissues under investigation. In carp muscle, MUFA predominated because it accounted for nearly half of the total fatty acids: 51.8% in TAG, followed by SFA: 29.3% and PUFA: 19.0%. PUFA represented the majority of the PL 54.8%, followed by SFA: 26.6% and

MUFA: 18.5%, similar to that reported in [25]. Carp muscle tissue contained only negligible amounts of F-acids, less than 0.05%. The composition of PL in male and female gonad tissues was found to be comparable to that of muscle PL (a high amount of PUFA, followed by SFA and MUFA), which was previously observed.

It was discovered that the triacylglycerol fraction of male gonad tissue contained the highest concentration of F-acids. F-acids were found in polar lipids and diacylglycerols in female gonad tissue, albeit at a significantly lower concentration than in male gonad tissue (Chvalova and Spicka, 2016).

This study aimed to develop a method for determining the fatty acids in fish roe and screening the FA profile of some common commercial fish roe.

## MATERIALS AND METHODS

All the solvents (*n*-hexane, *n*-heptane, methanol, sulphuric acid) were of analytical grade and chromatographic purity (acetonitrile), respectively, acquired from Merck KGaA, Darmstadt, Germany.

Commercial fish roes (tarama, cod, herring, lumpfish, salmon, and pike) from the supermarket were bought and used in 2018. The lipid phase was extracted with *n*-hexane, and the supernatant was derivatized according to the method published by Copolovici et al., 2017, to obtain the corresponding fatty acid methyl esters (Copolovici et al., 2017). These esters were identified by using GC-MS (Shimadzu 2010, Tokyo, Japan), mass spectrometry data from commercial libraries (NIST and Wiley) as presented in (Copolovici et al., 2017), and overlapping chromatograms are depicted in Figure 1.

## RESULTS AND DISCUSSIONS

Prior studies have shown high amounts of PUFA in fish roe, with some articles reporting levels as high as 50% of total FA (Al-Sayed Mahmoud et al., 2008; Intarasirisawat et al., 2011). Docosahexaenoic acid (C22:6 n-3) was found to be the most abundant fatty acid

(20.53–26.19%) in crude lipids extracted from tuna roes (51.22–54.90% of total lipids) (Al-Sayed Mahmoud et al., 2008). In our study, the values of PUFA are found to be from 26.61% in pike roe to 45.52% in cod roe (Table 1.)

In the present study, we found out that in the analyzed fish roes, the concentration of fatty acids varies depending on the species, so the myristic acid was found in a proportion of 3.55% in lumpfish roe and is increasing to up to 13.78% in pike roe. Elaidic acid was found in the samples analyzed, from 17.45% in pike roe to 26.26% in salmon roe (Figure 1).

As is revealed in Table 1, the fish roes analyzed present a good source of unsaturated fatty acids (UFA), with the following concentration for total  $\omega$ -3 FA: tarama < herring < pike < cod < salmon < lumpfish, while  $\omega$ -6 FA (linoleic acid) is found in a proportion of 6.46% in lumpfish roe and increase to 33.89% in tarama roe.




Figure 1. Overlapping chromatograms of salmon roe (blue), herring roe (pink), cod roe (black).

Table 1. Fatty acids identified in fish roe samples: the values are area expressed in % (R<sub>i</sub>: retention time).

Fatty acid	Molecular formula	Molecular weight (Da)	R <sub>i</sub> (min)	Tarama roe	Cod roe	Herring roe	Lumpfish roe	Salmon roe	Pike roe
Myristic acid (C14:0)	C <sub>14</sub> H <sub>28</sub> O <sub>2</sub>	228.371	5,407	5.70±0.04	5.85±0.03	10.48±0.14	3.58±0.11	3.67±0.31	13.78±0.28
Palmitoleic acid (C16:1, ω-7)	C <sub>16</sub> H <sub>30</sub> O <sub>2</sub>	254.408	9,726	0.95±0.03	1.35±0.17	1.58±0.15	4.64±0.15	6.36±0.16	5.30±0.21
Palmitic acid (C16:0)	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	256.424	10,552	16.48±0.17	16.30±0.42	19.65±0.21	13.62±0.02	11.45±0.14	22.50±0.35
Linoleic acid (C18:2, ω-6)	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	280.445	17,844	33.89±0.12	31.89±0.12	26.84±0.08	6.46±0.15	9.42±0.13	14.16±0.21
Elaidic acid(C18:1, ω-9)	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282.461	18,251	23.34±0.48	20.73±0.04	19.73±0.04	24.34±0.18	26.26±0.20	17.45±0.21
Oleic acid (C18:1, ω-7)	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282.461	18,49	1.52±0.02	2.95±0.02	1.95±0.02	5.19±0.16	4.73±0.37	3.23±0.14
Stearic acid (C18:0)	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	284.477	19,821	9.58±0.11	6.39±0.16	7.24±0.06	4.84±0.24	4.84±0.26	10.17±0.09
Eicosapentanoic acid (C20:5, ω-3)	C <sub>20</sub> H <sub>30</sub> O <sub>2</sub>	302.451	27,182	2.57±0.27	2.57±0.25	2.57±0.27	15.82±0.62	14.23±0.32	1.89±0.16
Docosahexaenoic acid (C22:6, ω-3)	C <sub>22</sub> H <sub>32</sub> O <sub>2</sub>	328.488	36,002	5.75±0.21	11.06±0.06	9.42±0.11	20.93±0.05	18.45±0.14	10.56±0.31
Σ SFA				31.76	28.54	37.37	22.00	19.96	46.45
Σ MUFA				25.81	25.02	23.245	34.16	37.34	25.98
Σ PUFA				42.20	45.52	38.83	43.20	42.09	26.61
Σ MUFA+ Σ PUFA				68.01	70.54	62.08	77.36	79.43	52.59
Σ SFA/ Σ UFA				0.47	0.40	0.60	0.28	0.25	0.88
Σ PUFA/ Σ MUFA				1.64	1.82	1.67	1.26	1.13	1.02

## CONCLUSIONS

Saturated fatty acids include palmitic acid (C16: 0) as the predominant FA in the analyzed samples, with a maximum of 22.5% in pike roe. The concentration of monounsaturated  $\omega$ -7 fatty acids (palmitoleic acid and oleic acid) varied between 2.47% (tarama roe) and 11.08% (salmon roe), while for  $\omega$ -9 fatty acid varied from 17.45% (pike roe) to 26.26% (salmon roe). Polyunsaturated fatty acids are smaller than 45.52%, while the concentration of  $\omega$ -3 fatty acids varied from 8.32% in tarama roe to 36.74% in lumpfish roe. The PUFA / MUFA ratio varies between 1.02% for pike roe and 1.82% for cod roe. Further studies are required to quantify the fatty acids from commercial fish roes and determine their correlation between chemical composition and biological activities.

## ACKNOWLEDGEMENTS

Funding for the equipment used in the study has been provided by The European Commission and The Romanian Government (project POS CCE 621/2014).

## REFERENCES

- Abdelhamid, A.S., Brown, T.J., Brainard, J.S., Biswas, P., Thorpe, G.C., Moore, H.J., Deane, K.H., Summerbell, C.D., Worthington, H.V., Song, F.J., Hooper, L., 2020. Omega-3 fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews*.
- Abdelhamid, A.S., Brown, T.J., Brainard, J.S., Biswas, P., Thorpe, G.C., Moore, H.J., Deane, K.H.O., AlAbdulghafoor, F.K., Summerbell, C.D., Worthington, H.V., Song, F.J., Hooper, L., 2018. Omega-3 fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews*, CD003177.
- Al-Sayed Mahmoud, K., Linder, M., Fanni, J., Parmentier, M., 2008. Characterisation of the lipid fractions obtained by proteolytic and chemical extractions from rainbow trout (*Oncorhynchus mykiss*) roe. *Process Biochemistry* 43, 376-383.
- Alagawany, M., Elnesr, S.S., Farag, M.R., Abd El-Hack, M.E., Khafaga, A.F., Taha, A.E., Tiwari, R., Yatoo, M.I., Bhatt, P., Khurana, S.K., Dhama, K., 2019. Omega-3 and Omega-6 Fatty Acids in Poultry Nutrition: Effect on Production Performance and Health. *Animals* 9.
- AlAmmar, W.A., Albeesh, F.H., Ibrahim, L.M., Algindan, Y.Y., Yamani, L.Z., Khattab, R.Y., 2021. Effect of omega-3 fatty acids and fish oil supplementation on multiple sclerosis: a systematic review. *Nutritional Neuroscience* 24, 569-579.
- Berger, M.E., Smesny, S., Kim, S.W., Davey, C.G., Rice, S., Sarnyai, Z., Schlogelhofer, M., Schafer, M.R., Berk, M., McGorry, P.D., Amminger, G.P., 2017. Omega-6 to omega-3 polyunsaturated fatty acid ratio and subsequent mood disorders in young people with at-risk mental states: a 7-year longitudinal study. *Translational Psychiatry* 7.
- Chvalova, D., Spicka, J., 2016. Identification of furan fatty acids in the lipids of common carp (*Cyprinus carpio* L.). *Food Chemistry* 200, 183-188.
- Copolovici, D., Bungau, S., Boscencu, R., Tit, D.M., Copolovici, L., 2017. The Fatty Acids Composition and Antioxidant Activity of Walnut Cold Press Oil. *Revista De Chimie* 68, 507-509.
- Cretton, M., Malanga, G., Sobczuk, T.M., Mazzuca, M., Marine lipids as a source of high-quality fatty acids and antioxidants. *Food Reviews International*.
- Dawczynski, C., Dittrich, M., Neumann, T., Goetze, K., Welzel, A., Oelzner, P., Volker, S., Schaible, A.M., Troisi, F., Thomas, L., Pace, S., Koeberle, A., Werz, O., Schlattmann, P., Lorkowski, S., Jahreis, G., 2018. Docosahexaenoic acid in the treatment of rheumatoid arthritis: A double-blind, placebo-controlled, randomized cross-over study with microalgae vs. sunflower oil. *Clinical Nutrition* 37, 494-504.
- de Castro, G.S., Andrade, M.F., Pinto, F.C.S., Faiad, J.Z., Seelaender, M., 2022. Omega-3 Fatty Acid Supplementation and Its Impact on Systemic Inflammation and Body Weight in Patients With Cancer Cachexia-A Systematic Review and Meta-Analysis. *Frontiers in Nutrition* 8.
- de la Guia-galipienso, F., Martinez-Ferran, M., Vallecillo, N., Lavie, C.J., Sanchis-Gomar, F.,

Pareja-Galeano, H., 2021. Vitamin D and cardiovascular health. *Clinical Nutrition* 40, 2946-2957.

Franchi, C., Ardoino, I., Bosetti, C., Negri, E., Serraino, D., Crispo, A., Giacosa, A., Fattore, E., Dolci, A., Bravi, F., Turati, F., La Vecchia, C., D'Avanzo, B., 2022. Inverse Association between Canned Fish Consumption and Colorectal Cancer Risk: Analysis of Two Large Case-Control Studies. *Nutrients* 14.

Hansen, A.L., Ambroziak, G., Thornton, D., Dahl, L., Molde, H., Grung, B., 2020. Fatty Fish Intervention and Psychophysiological Responses to Mental Workload in Forensic Inpatients A Randomized Control Trial. *Journal of Psychophysiology* 34, 10-18.

Hsieh, S.L., Li, J.H., Dong, C.D., Chen, C.W., Wu, C.C., 2022. Carnosine suppresses human colorectal cancer cell proliferation by inducing necroptosis and autophagy and reducing angiogenesis. *Oncology Letters* 23.

Intarasirisawat, R., Benjakul, S., Visessanguan, W., 2011. Chemical compositions of the roes from skipjack, tongol and bonito. *Food Chemistry* 124, 1328-1334.

Jiang, W., Li, F.R., Yang, H.H., Chen, G.C., Hua, Y.F., 2021. Relationship Between Fish Oil Use and Incidence of Primary Liver Cancer: Findings From a Population-Based Prospective Cohort Study. *Frontiers in Nutrition* 8.

Laumann, R.D., Iversen, T., Frandsen, T.L., Molgaard, C., Stark, K.D., Schmiegelow, K., Lauritzen, L., 2022. Whole blood long-chain n-3 fatty acids as a measure of fish oil compliance in children with acute lymphoblastic leukemia: a pilot study. *Prostaglandins Leukotrienes and Essential Fatty Acids* 177.

Lofvenborg, J.E., Andersson, T., Carlsson, P.O., Dorkhan, M., Groop, L., Martinell, M., Tuomi, T., Wolk, A., Carlsson, S., 2014. Fatty fish consumption and risk of latent autoimmune diabetes in adults. *Nutrition & Diabetes* 4.

Sikka, P., Behl, T., Sharma, S., Sehgal, A., Bhatia, S., Al-Harrasi, A., Singh, S., Sharma, N., Aleya, L., 2021. Exploring the therapeutic potential of omega-3 fatty acids in depression. *Environmental Science and Pollution Research* 28, 43021-43034.

Stasiewicz, B., Wadolowska, L., Biernacki, M., Slowinska, M.A., Stachowska, E., 2022. Dietary Fat Intake: Associations with Dietary Patterns and Postmenopausal Breast Cancer-A Case-Control Study. *Cancers* 14.

Wang, Y., Liu, K., Long, T.T., Long, J.Y., Li, Y., Li, J.Y., Cheng, L.M., 2022. Dietary fish and omega-3 polyunsaturated fatty acids intake and cancer survival: A systematic review and meta-analysis. *Critical Reviews in Food Science and Nutrition*.

Yonezawa, K., Kusumoto, Y., Kanchi, N., Kinoshita, H., Kanegae, S., Yamaguchi, N., Ozawa, H., 2020. Recent trends in mental illness and omega-3 fatty acids. *Journal of Neural Transmission* 127, 1491-1499.

Zhang, T., Wu, S.S., Xu, F.Y., Chang, J.P., Guo, Y.X., Zhou, Z.T., Xu, R.W., Wang, T.C., Wang, X.M., Wang, M.H., Qin, S.Y., Zhang, S.G., Li, J., 2022. The Association between Dietary Protein Intake and the Risk of Pancreatic Cancer: Evidence from 14 Publications. *Nutrition and Cancer-an International Journal*.

ISSN 1582-1021

e-ISSN 2668-4764

Edited by "AUREL VLAICU" University  
Arad, Romania



Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.